

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : G01N 33/53, 33/577, 33/543, A61K 35/00, 31/19, 31/195, 31/66	A1	(11) International Publication Number: WO 97/42501 (43) International Publication Date: 13 November 1997 (13.11.97)
---	-----------	---

(21) International Application Number: **PCT/US97/07127**
(22) International Filing Date: **7 May 1997 (07.05.97)**

(30) Priority Data:
60/016,322 7 May 1996 (07.05.96) **US**

(71) Applicant: **BOEHRINGER INGELHEIM PHARMACEUTI-
CALS, INC. [US/US]; 900 Ridgebury Road, P.O. Box 368,
Ridgefield, CT 06877 (US).**

(72) Inventors: **BARTON, Randall, W.; 10 Paul Spring Hollow,
Farmington, CT 06032 (US). KENNEDY, Charles, A.; 99
East Side Boulevard, Waterbury, CT 06706 (US).**

(74) Agents: **RAYMOND, Robert et al.; Boehringer Ingelheim Cor-
poration, 900 Ridgebury Road, P.O. Box 368, Ridgefield,
CT 06877 (US).**

(81) Designated States: **CA, JP, MX, European patent (AT, BE,
CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
PT, SE).**

Published

*With international search report.
Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: **PHOSPHOTYROSINE MIMICS AND METHODS FOR IDENTIFYING AND USING SAME**

(57) Abstract

This invention relates to phosphotyrosine mimics which, when incorporated into an appropriate molecular structure, are capable of inhibiting the binding of tyrosine kinase-dependent regulatory proteins to the native phosphotyrosine-containing receptors. This invention also relates to methods and kits for identifying and using phosphotyrosine mimics. The compounds, kits and methods of this invention are useful for identifying and designing antagonists of tyrosine-kinase dependent regulatory proteins, such as signal transduction proteins containing SH₂ binding domains.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Phosphotyrosine Mimics and Methods for Identifying And Using Same

Technical Field of Invention

5 This invention relates to phosphotyrosine mimics which, when incorporated into an appropriate molecular structure, are capable of inhibiting the binding of tyrosine kinase-dependent regulatory proteins to their native phosphotyrosine - containing ligands or receptors. This invention also relates to methods and kits for identifying
10 and using phosphotyrosine mimics. The compounds, kits and methods of this invention are useful for identifying and designing antagonists of tyrosine-kinase dependent regulatory proteins, such as signal transduction proteins containing SH₂ binding domains

Background of the Invention

15 The activation of cells by growth factors, mitogens or other cytokines to undergo proliferation and/or differentiation is often dependent on inducible tyrosine kinase activity. This tyrosine kinase activity increases the phosphotyrosine content of many receptor-like and cytoplasmic regulatory proteins. Often, the physical association of
20 such regulatory proteins is mediated through those phosphotyrosine residues.

For example, one mechanism of cellular regulation involves the physical association of signal transduction proteins with one or more phosphorylated tyrosine-containing receptor subunits, called immunoreceptor tyrosine-based activation motifs (ITAMs),
25 present on its native ligand or receptor. This association is a common feature of many cytoplasmic signal transduction pathways, as well as other immunologically important regulatory protein-receptor based interactions (M.A. Osborne et al., BioTechnology, 13 (1995)). Of particular importance is the interaction between
30 phosphorylated ITAMs and regulatory proteins containing Src homology domain 2 regions ("SH₂ binding domains"). Examples of immunologically important proteins containing SH₂ binding domains include ZAP-70, Fyn, Lyn, Lnk, Abl, Vav, Huk, Blk, PLC γ 1, GAP, Crk, Shc and p⁵⁶lck. Although proteins having SH₂ binding domains tend to have sequence-specific affinities for their ITAM containing ligands

or receptors, the binding interaction itself is ubiquitously mediated through one or more phosphorylated tyrosine residues. Therefore, the presence of phosphorylated tyrosine plays a critical role in signal transduction involving virtually all proteins containing SH₂ binding domains.

5

Given the above understanding of signal transduction and cellular regulation, it follows that growth factor- and cytokine-induced cell proliferation and/or differentiation can be selectively inhibited by antagonizing the interaction of regulatory proteins dependent on tyrosine kinase activity with their native phosphotyrosine-containing ligands or receptors. Such antagonists would undoubtedly be useful to treat a variety of disorders, including those associated with or caused by neoplastic diseases or chronic inflammatory diseases.

To date, however, antagonists of tyrosine kinase-dependent regulatory proteins have not fulfilled their potential as useful pharmaceutical agents. One major hurdle has been the necessary inclusion of a phosphorylated α -amino acid residue or a phosphorylated analog thereof, to perform the crucial role of the native tyrosine phosphoprotein ligands or receptors of these regulatory proteins. However, agents containing phosphotyrosine, other phosphorylated α -amino acid residues, or phosphorylated analogs thereof, cannot generally be used as therapeutic agents because the presence of the phosphorylated moiety substantially impedes cell penetrability. Until now, no effective replacement or mimic for the critical phosphotyrosine residue has been identified. Further, no accurate methods for identifying such phosphotyrosine mimics have existed. Accordingly, the need exists for effective phosphotyrosine mimics and convenient methods for identifying such mimics.

Brief Description of the Invention

This invention satisfies the above-mentioned needs by providing phosphotyrosine mimics and convenient methods for identifying phosphotyrosine mimics.

One object of this invention is to provide a method for identifying phosphotyrosine mimics comprising the steps of:

- (a) contacting a non-phosphorylated test compound and a phosphotyrosine-containing substance with an anti-phosphotyrosine antibody; and
- (b) detecting the ability of the non-phosphorylated test compound to bind to the anti-phosphotyrosine antibody.

Another object of this invention is to provide phosphotyrosine mimics that, when incorporated into an appropriate molecular structure, inhibit the binding of tyrosine kinase-dependent regulatory proteins to their native phosphotyrosine-containing ligands or receptors.

Yet another object of this invention is to provide kits for determining the presence of a phosphotyrosine mimic in a non-phosphorylated test compound.

10 A further object of this invention is to provide methods for using phosphotyrosine mimics to inhibit the activation of tyrosine kinase-dependent regulatory proteins.

Yet a further object of this invention is to provide a method for treating neoplastic or chronic inflammatory disease using the phosphotyrosine-containing compounds according to this invention.

Other objects and advantages of this invention will be apparent to those of ordinary skill in the art.

20 Detailed Description of the Invention

Unless otherwise indicated, the following definitions apply:

25 The terms "amino acid" and " α -amino acid" are used interchangeably herein and refer to the naturally occurring α -amino acids, as well as those amino acids in their D-configurations and non-native, synthetic and modified amino acids known to those of ordinary skill in the art (e.g., homocysteine, ornithine, norleucine and β -valine).

30 The term "patient" refers to a warm-blooded animal, such as a human, who is afflicted with a neoplastic or chronic inflammatory disorder.

35 The term "phosphotyrosine mimic" refers to a non-phosphorylated chemical moiety which is functionally capable of replacing phosphotyrosine in a native phosphotyrosine-containing ligand. Specifically, when incorporated into an appropriate molecular structure, a phosphotyrosine mimic is capable of antagonizing the binding of a tyrosine kinase dependent regulatory protein to its natural ligand(s). The antagonistic ability of such compounds may be detected by any of the detection methods described herein, or any other conventional detection method known to those

of ordinary skill in the art. Preferably, a phosphotyrosine mimic-containing compound according to this invention is capable of inhibiting the binding of a tyrosine kinase-dependent regulatory protein to its corresponding, phosphorylated ligand or receptor by at least about 30% at a concentration of 10 μ M (more preferably, by at least about 35%, even more preferably, by at least about 40%, even more preferably, by at least about 50% and most preferably, by at least 60%, 70% or even 80%).

Preferred phosphotyrosine mimic-containing compounds according to this invention are characterized by improved cell penetrability over that possessed by the corresponding phosphotyrosine-containing analogs. The improved cell penetrability possessed by these compounds advantageously allows the compounds to more readily pass through the cell membrane, thereby increasing the likelihood of physical interaction with the targeted regulatory protein (such as a particular targeted phosphotyrosine kinase-dependent regulatory protein containing an SH₂ binding domain).

The present invention also provides a convenient and sensitive method for identifying phosphotyrosine mimics by utilizing anti-phosphotyrosine antibodies to rapidly screen large numbers of non-phosphorylated test compounds. Specifically, this method comprises the steps of:

- (a) contacting a non-phosphorylated test compound and a phosphotyrosine-containing substance with an anti-phosphotyrosine antibody; and
- (b) detecting the ability of the non-phosphorylated test compound to inhibit the binding of the anti-phosphotyrosine antibody to the phosphotyrosine-containing substance.

In a preferred embodiment, the method for identifying a phosphotyrosine mimic according to this invention comprises the steps of :

- (a) competitively binding a non-phosphorylated test compound and a phosphotyrosine-containing substance with an anti-phosphotyrosine antibody; and
- (b) detecting the degree to which the non-phosphorylated test compound displaces the phosphotyrosine-containing substance.

More preferably, the method for identifying a phosphotyrosine mimic according to this invention comprises the steps of :

- (a) incubating a non-phosphorylated test compound with an anti-phosphotyrosine antibody to form a solution;

- (b) contacting the solution of step (a) with a known amount of a phosphotyrosine-containing substance; and
- (c) detecting the degree to which the non-phosphorylated test compound displaces the phosphotyrosine-containing substance.

5

In each of the above embodiments, preferred non-phosphorylated test compounds containing a phosphotyrosine mimic according to this invention are capable of inhibiting the binding of the anti-phosphotyrosine antibody to the phosphotyrosine-containing substance by at least about 30% at a concentration of 10 μ M (more preferably, by at least about 35%, even more preferably, by at least about 40%, even more preferably, by at least about 50% and most preferably, by at least 60%, 70% or 80%).

The anti-phosphotyrosine antibodies useful in the methods of this invention are those which are capable of binding to a phosphotyrosine residue present in a phosphotyrosine-containing compound, wherein the phosphotyrosine residue is free of steric hindrance or other blockage to reaction with the antibody. Such anti-phosphotyrosine antibodies are well known to those of ordinary skill in the art and may be easily obtained by in-vivo production and harvesting according to known techniques. In general, conventional techniques for producing and harvesting monoclonal anti-phosphotyrosine antibodies comprise injecting a mammal (e.g. a mouse) with a compound containing a phosphotyrosine residue, fusing the mammal's spleen cells with myeloma cells and assaying the resultant hybridoma cells for antibodies which specifically bind to phosphotyrosine residues in different phosphotyrosine-containing proteins containing phosphotyrosine. Once identified, the cells producing these antibodies can be cloned and grown in mass culture or injected into an animal (e.g., a mouse) to induce secretion of the desired antibody. As those of ordinary skill in the art will readily appreciate, other conventional methods for producing anti-phosphotyrosine antibodies may be used equally well for the purposes of this invention.

Alternatively, anti-phosphotyrosine antibodies may be purchased commercially (for example, PY-20, 4G10, B4, RC20, 1G2, 25.2G4, 3-365-10, 6D12, PT66, PY54, PY69 and Z027 are available from suppliers such as Transduction Labs (Lexington, KY), Zymed (South San Francisco, CA), Upstate Biotechnology (Lake Placid, NY), Biogenesis LTD (Sandown, NH), Boehringer Mannheim Biochemicals (Indianapolis, IN), Wako Chemicals USA Inc (Richmond, VA), Medical & Biological Labs LTD (Nagoya, Japan), Sigma Chemical Co (St Louis MO), Biodesign International

(Kennebunk ME), and ICN Biochemicals (Costa Mesa, CA) . Although the methods of this invention envision use of any suitable anti-phosphotyrosine antibody (e.g., polyclonal or monoclonal; human or non-human), preferably the antibody is a commercially available, monoclonal antibody which specifically and selectively
5 targets the phosphotyrosine residue of one or more naturally occurring phosphotyrosine-containing ligands. More preferably, the antibody is a mouse anti-phosphotyrosine monoclonal antibody. Most preferably the antibody is PY-20.

The methods of this invention employ assays which measure anti-phosphotyrosine
10 antibody binding affinity of a non-phosphorylated test compound versus a phosphotyrosine-containing substance. Although all assays that perform this function are contemplated by this invention, direct binding assays are less preferred because they tend to be difficult to carry out in a high-throughput manner. Competitive
15 binding assays are preferred because they are more easily and quickly performed with a large number of non-phosphorylated test compounds. Many such competitive binding assays are well known to those of ordinary skill in the art. Competitive binding assays which may be used in the methods of this invention are typically immunoassays that detect the presence or concentration of the free or bound forms of anti-phosphotyrosine antibody, phosphotyrosine-containing substance or non-
20 phosphorylated test compound. Depending on precisely which immunoassay is used, the free- and bound-forms of those substances may be readily distinguished using an appropriate label.

Typically, the known phosphotyrosine-containing substance or the anti-
25 phosphotyrosine antibody for use in the methods of this invention is conjugated with a conventional monitorable label which does not interfere with the ability of the known phosphotyrosine-containing substance and the anti-phosphotyrosine antibody to cross-react. Labels that are generally useful for this purpose include, but are not limited to, chemical ligands, enzymatic labels, radioactive labels, fluorescent labels,
30 luminescent labels and electronic labels. One particularly useful chemical ligand is biotin. Particularly useful enzymatic labels include, but are not limited to, horseradish peroxidase, β -galactosidase, alkaline phosphatase and acetylcholinesterase. Particularly useful radioactive ligands include, but are not limited to, phosphorus ^{32}P and ^{33}P , iodine ^{125}I and ^{131}I and hydrogen, ^3H . Particularly useful fluorescent
35 labels include, but are not limited to, rhodamine, tetramethyl rhodamine isothiocyanate, phycoerythrin and dansyl chloride. One particularly useful luminescent label is luminol.

Particularly preferred competitive binding immunoassays useful in the methods of this invention include, but are not limited to, enzyme-linked, fluorescent, chemiluminescent, radio and biosensor immunoassays. Although the immunoassays of this invention may be carried out in solution or on a solid support, we prefer using
5 a solid support (such as the wells of a microtiter plate) to facilitate large-scale screening of non-phosphorylated test compounds. Preferably, the immunoassay used in the methods of this invention is ELISA, a fluorescent immunoassay or a biosensor immunoassay.

10 In an enzyme-linked immunosorbent assay (ELISA) according to this invention, the known phosphotyrosine-containing substance or the anti-phosphotyrosine antibody may be directly labeled with an enzyme or indirectly labeled with an enzyme-labeled antibody. The enzyme labels useful for ELISA are those that, under appropriate conditions, catalyze an observable reaction with a given enzymatic substrate. Such
15 enzymatic activity is typically measured by formation of a colored or otherwise easily identifiable reaction product. In a typical ELISA scheme according to this invention, the unlabeled compound (either the known phosphotyrosine-containing substance or the anti-phosphotyrosine antibody) is bound to a solid support and the correspondingly labeled substrate is added. After any unbound species are washed
20 away, the non-phosphorylated test compound is added. Unbound species are again removed by washing and the enzyme is activated to determine the extent to which the test compound displaced the known phosphotyrosine-containing substance. Wherever the phosphotyrosine-containing substance was displaced, no enzymatic activity will be detected.

25 Fluorescent immunoassays useful in the methods of this invention involve the use of conventional fluorochromes. Such fluorochromes may be linked directly to the anti-phosphotyrosine antibody or the known ligand used in the immunoassay, or alternatively, the fluorochrome may be linked indirectly to those compounds with
30 fluorochrome-labeled antibodies. Fluorochromes are dyes that absorb radiation (such as UV light) and emit light of a different characteristic frequency when the labeled compound is bound to a substrate or present in its free state. Therefore, when a known phosphotyrosine-containing substance according to this invention is labeled with a fluorochrome, and the anti-phosphotyrosine antibody and non-phosphorylated
35 test compound are subsequently added, it is a routine matter to detect the extent to which the known phosphotyrosine-containing substance has been displaced by the test compound (e.g., by measuring the amount of free, known phosphotyrosine-containing substance).

In biosensor immunoassays, the binding of an anti-phosphotyrosine antibody to a known phosphotyrosine-containing substance is detected by measuring the change in refractive index that occurs when the solution-phase antibody binds to the known phosphotyrosine-containing substance which has been tethered to an optically sensitive surface. The biosensor allows the determination of equilibrium binding constants together with on and off rates of the interacting molecules. Therefore, when a known phosphotyrosine-containing substance is bound to an optically sensitive surface and the anti-phosphotyrosine antibody and non-phosphorylated test compound are subsequently added, it is a routine matter to detect the extent to which the binding of the anti-phosphotyrosine antibody to the known phosphotyrosine-containing substance has been displaced by the test compound.

This invention also provides a kit for the rapid detection of phosphotyrosine mimics comprising:

- (a) an immobilized phosphotyrosine-containing substance and
- (b) an anti-phosphotyrosine antibody, wherein said antibody is conjugated to a detectable label.

Alternatively, the kit comprises:

- (a) an immobilized anti-phosphotyrosine antibody and
- (b) a phosphotyrosine-containing substance, wherein said substance is conjugated to a detectable label.

Preferred kits are those that comprise the preferred phosphotyrosine-containing substances and preferred anti-phosphotyrosine antibody described above for use in the methods of this invention. The kits of this invention may also include buffers, standards and other conventional reagents to facilitate their use. Preferably, the immobilized component of the kit is covalently bound to a solid resin (such as microtiter plate). Techniques for producing these kits will be readily apparent to those of ordinary skill in the art.

Using the methods and kits of this invention, a panel of non-phosphorylated test compounds may be quickly and efficiently assayed for the presence of a phosphotyrosine mimic. Once identified, these phosphotyrosine mimics may be incorporated into an appropriate molecular structure to provide compounds which inhibit the binding of tyrosine kinase-dependent regulatory proteins to their native

receptors. Accordingly, this invention also provides a method for inhibiting the activation of a tyrosine kinase-dependent regulatory protein comprising the steps of:

- (a) incorporating a phosphotyrosine mimic according to this invention into an appropriate molecular structure to produce a phosphotyrosine mimic containing
5 compound, and
- (b) contacting the compound of step (a) with a tyrosine-kinase dependent regulatory protein.

Preferably, the compound of step (a) is covalently attached to a solid support prior to
10 the binding step (b) using known techniques (such as those useful for the kits of this invention).

An example of an appropriate molecular structure for the above detailed use is a phosphotyrosine-containing peptidomimetic in which the phosphotyrosine can be
15 replaced with a phosphotyrosine mimic. Preferably, the peptidomimetic comprises a fragment of a native tyrosine kinase-dependent regulatory protein ligand or receptor, in which the naturally occurring phosphotyrosine residue is replaced with a phosphotyrosine mimic identified by the methods of this invention. Such peptidomimetics generally include from about 1 to about 30 of the naturally occurring
20 α -amino acid residues flanking each side of the naturally occurring phosphotyrosine residue. More preferably, the peptidomimetic comprises from about 1 to about 10 (and most preferably, from about 1 to about 5) of the naturally occurring α -amino acid residues flanking each side of the naturally occurring phosphotyrosine residue. Although the naturally occurring α -amino acids are preferred in these
25 peptidomimetics, those naturally occurring α -amino acids may be optionally modified or substituted according to known techniques. Preferred modifications and substitutions to the naturally occurring amino acid sequence are conservative ones (i.e., those having a minimal influence on the secondary structure and hydrophobic nature of the peptide). These include those substitutions and modification described
30 in Dayhoff, Atlas of Protein Sequence and Structure, 5 (1978) and Argos, EMBO J., 8, pp. 779-85 (1989).

Peptidomimetics according to this invention may be prepared using known techniques. In general, a phosphotyrosine mimic is used to replace phosphotyrosine
35 in the native peptide sequence by first synthesizing a modified residue containing the phosphotyrosine mimic (having both a carboxy and amino terminus) and using this modified residue to synthesize the peptidomimetic using a known schemes for peptide synthesis.

The above described peptidomimetics may be prepared using any conventional peptide production methodology including solid phase or solution phase synthesis, recombinant DNA technology and combinations thereof. Preferably, these

5 peptidomimetics are produced using solid phase synthesis. The solid support may be any suitable resin conventionally employed in the art. Preferred resins include, but are not limited to p-benzyloxyalcohol polystyrene and p-methylbenzhydrylamine. Amino acids for use in this method may be side chain protected, if necessary. The

10 criteria for choosing an appropriate side chain protecting group include: (a) stability of the side chain protecting group to the reaction conditions needed for removal of the α -amino protecting group, (b) stability of the side chain protecting group to the reaction conditions required for amino acid coupling and (c) removability of the side chain protecting group upon the conclusion of peptide synthesis and under conditions

15 that do not otherwise effect the peptide structure. The first amino acid is amino protected then coupled to the resin. Amino protecting groups include, but are not limited to, 9-fluorenyl-methyloxycarbonyl (Fmoc) and t-butoxycarbonyl (BOC). The amino protecting group is then removed using conventional methods. After removal of the amino protecting group, the remaining amino-protected amino acids (side-chain protected, if necessary) are sequentially added to produce the desired

20 peptidomimetic.

The phosphotyrosine mimic identified according to this invention may also be functional group protected, if desired, and added to the growing peptide chain as described above. Functional group protection is well within the ordinary skill of the

25 art and is typically carried out as described above for side group protection. Other protecting groups useful for side chain or functional group protection according to this invention may be found in well known organic chemistry references. It should also be appreciated that phosphotyrosine mimic-containing compounds according to this invention include the free form of such compounds as well as pharmaceutically

30 acceptable salts of those compounds, where such forms exists.

Using the above methodology, phenyl carboxylic acid was identified as a phosphotyrosine mimic according to this invention. Although unsubstituted phenyl carboxylic acid is the preferred phosphotyrosine mimic, this invention also envisions

35 the use of substituted-phenyl analogs of phenyl carboxylic acid (referred to collectively herein as phenyl carboxylic acid mimics). Preferred phenyl substituents include a member of the group consisting of halogens (especially, chloride, fluoride and bromide), branched or unbranched C₁₋₅ alkyl, C₁₋₅ alkoxy, C₂₋₅ alkenyl or

- alkynyl, (wherein any one of the foregoing alkyl, alkoxy, alkenyl or alkynyl groups may be unsubstituted or substituted with cyano, nitro, amino, trifluoromethyl, trifluoromethoxy or hydroxy and wherein any unsubstituted methylene may be replaced with O, S or C=O), cyano, nitro, amino, trifluoromethyl, trifluoromethoxy and hydroxy. The phenyl carboxylic acid mimic may replace a phosphotyrosine residue in an appropriate molecular structure by attachment at any unsubstituted position on the phenyl ring. Preferably, the phenyl ring is attached to the molecular structure at the 2-position.
- 5
- 10 The phenyl carboxylic acid mimics according to this invention are structurally simple and advantageously, easily synthesized from commercially available starting materials using conventional synthetic techniques. Similarly, incorporation of a phenyl carboxylic acid mimic into an appropriate molecular structure (and preferably, a peptidomimetic of a native tyrosine kinase-dependent regulatory protein ligand or
- 15 receptor) is readily accomplished using known techniques and, preferably, the methodology detailed above. Specifically, the phenyl carboxylic acid phosphotyrosine mimic may be used to replace phosphotyrosine in a peptidomimetic by first synthesizing an α -amino acid residue containing the phosphotyrosine mimic (having both a carboxy and amino terminus) and using this modified residue to
- 20 synthesize the peptidomimetic using any known scheme for peptide synthesis.

It should be understood that only those compounds having combinations of variables that result in stable structures are included within the scope of this invention. Stable structures are those that can be produced according to the above mentioned techniques and stored for an acceptable period of time. Preferably, the stable structures are those that can be stored at 32°F (0°C) for at least one week without detectable levels of decomposition. In addition to the free forms of these compounds, this invention also includes the pharmaceutically acceptable salts thereof. The production of pharmaceutically acceptable salts of compounds according to this invention is well

25

30 within the ordinary skill in the art.

Without wishing to be bound by theory, we believe that the phosphotyrosine mimics according to this invention, when incorporated into an appropriate molecular structure, are capable of binding tyrosine-kinase dependent regulatory proteins in a manner substantially similar or identical to that of the native phosphotyrosine-containing receptor. Therefore, the phosphotyrosine mimic-containing compounds are competitive inhibitors of that binding. Specifically, once bound, the phosphotyrosine mimics inhibit the ability of the regulatory protein to bind its native

35

phosphotyrosine-containing receptor, thereby inhibiting cellular activation. It will be apparent from the nature of this discovery that the immunoassay methods and kits of this invention may be advantageously used to identify phosphotyrosine mimics without reference to a particular tyrosine kinase-dependent regulatory protein. In contrast to mimics that might be identified by competitive binding assays to a specified tyrosine kinase-dependent regulatory protein, the phosphotyrosine mimics identified according to this invention possess a more general utility. By incorporating these mimics into an appropriate molecular structure (as described above), the phosphotyrosine mimics according to this invention may be integrally used as antagonists to virtually any tyrosine kinase-dependent regulatory protein.

As antagonists of tyrosine kinase-dependent regulatory proteins, the phosphotyrosine mimic-containing structures according to this invention can be used to treat a variety of disorders, including those associated with or caused by neoplastic diseases or chronic inflammatory diseases. Specifically, by blocking or displacing the binding of the native phosphotyrosine-containing ligands or receptors of these regulatory proteins, the phosphotyrosine mimic-containing structures according to this invention effectively disrupt the associated regulatory cascades. Representative neoplastic diseases that are treatable with the phosphotyrosine mimic-containing antagonists according to this invention include (but are not limited to): leukemias (including, but not limited to, acute lymphocytic, acute lymphoblastic, chronic lymphocytic, acute myeloblastic and chronic myelocytic), carcinomas (including, but not limited to, adenocarcinoma and that of the colon, ovaries, cervix, esophagus, stomach, small intestines, pancreas and lungs), sarcomas (including, but not limited to, osteosarcoma, leiomyoma, liposarcoma, hemangioma, hemangiosarcoma and Kaposi's sarcoma), malignant melanomas (including, but not limited to, amelanotic and melanotic), mixed types of neoplasias (such as, but not limited to, carcinosarcoma, lymphoid tissue type, follicular reticulum, cell sarcoma and Hodgkin's disease), neuroblastoma, cerebral malaria, capillary leak syndrome, hematological malignancies and the like. Representative chronic inflammatory diseases treatable with the phosphotyrosine mimic-containing antagonists according to this invention include (but are not limited to): rheumatoid arthritis, multiple sclerosis, Guillain-Barre syndrome, Crohn's disease, ulcerative colitis, psoriasis, graft versus host disease, lupus erythematosus and insulin-dependent diabetes mellitus.

The present invention provides pharmaceutical compositions comprising a therapeutically effective amount of a phosphotyrosine mimic-containing substance according to this invention or, if desired, a pharmaceutically acceptable salt thereof.

The compounds or pharmaceutical compositions according to this invention may be used to treat any of the disease states mentioned above by administering a therapeutically effective amount of a compound or composition according to this invention to a patient. For such treatment, the preferred phosphotyrosine mimic-
5 containing substance is a peptidomimetic of a phosphotyrosine-containing ligand or receptor of a tyrosine-kinase dependent regulatory protein. Preferably, the phosphotyrosine mimic is phenyl carboxylic acid or a functional group protected derivative thereof.

10 A therapeutically effective amount refers to an amount which is effective, upon single or multiple dose administration to the patient, in controlling the growth of a neoplasm, in alleviating, in whole or in part, the symptoms of the chronic inflammatory disorder, prolonging the survivability or improving the clinical disposition or physical well-being of the patient. Treatment according to this
15 invention may or may not completely eradicate the symptoms or the disorder being treated. A therapeutically effective amount can be readily determined by the attending diagnostician by the use of known techniques and by observing the results obtained under analogous circumstances. In determining a therapeutically effective amount, a number of factors are considered by the attending diagnostician, including
20 (but not limited to): the species of mammal; its size, age and general health; the specific disease involved; the degree of or involvement or the severity of the disease; the response of the individual patient; the particular compound or composition administered; the mode of administration; the bioavailability characteristics of the preparation administered; the dose regimen selected; the use of concomitant
25 mediation and other relevant circumstances.

The compounds and pharmaceutical compositions according to this invention may be administered to the patient in any pharmaceutically acceptable and effective dosage form. Examples of such dosage forms include (but are not limited to): intravenous,
30 intramuscular, subcutaneous, intra-articular, intrasynovial, intrathecal, periosteal, intratumoral, peritumoral, intralesional, perilesional, infusion, sublingual, buccal, transdermal, oral, topical or inhalation. Preferred dosage forms include oral, topical, intravenous, subcutaneous and transdermal.

35 In general, a therapeutically effective amount of a compound according to this invention is expected to vary in the range of about 0.1 mg/kg body weight/day (mg/kg/day) to about 100 mg/kg/day. Preferred amounts should range from about 0.1 mg/kg/day to about 50 mg/kg/day. More specifically, preferred dosage levels for

various modes of administration are: intravenous (from about 0.1 mg/kg/day to about 40 mg/kg/day); intramuscular (from about 1 mg/kg/day to about 50 mg/kg/day); orally (from about 5 mg/kg/day to about 100 mg/kg/day); intranasal instillation (from about 5 mg/kg/day to about 100 mg/kg/day); and aerosol (from about 5 mg/kg/day to about 100 mg/kg/day).

Dosage forms may include any pharmaceutically acceptable carriers and adjuvants that are known to those of ordinary skill in the art. These carriers and adjuvants include, for example, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins (such as human serum albumin), buffer substances (such as phosphates), glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes (such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate or sodium chloride), zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances and glycols (such as polyethylene glycol). Such forms include (but are not limited to) tablet, capsule, caplet, liquid, solution, suspension, emulsion, lozenges, syrup, reconstitutable powder, granule, suppository and transdermal patch. Methods for preparing such dosage forms are well known (see, for example, H.C. Ansel and N.G. Popovich, Pharmaceutical Dosage Forms and Drug Delivery Systems, 5th edition, Lea and Febiger, 1990).

The compounds of this invention may be administered alone or in combination with one or more conventional anti-neoplastic or anti-inflammatory agent. Such agents are well known and may be selected as circumstances dictate by those of ordinary skill in the art. Advantageously, such combination therapy may utilize lower dosages of the conventional therapeutic agents, thereby avoiding possible toxicity and adverse side effects incurred when those agents are administered as monotherapies. For example, the compounds of this invention may be used in combination with conventional cancer drugs (such as methotrexate, taxol, 5-fluorouracil, cis-platinum, cortisone, nitrogen mustards, thiotepa and nitrosoureas) and conventional anti-inflammatory drugs (such as non-steroidal anti-inflammatory agents, penicillamine, methotrexate, cortisone and gold salts).

Besides their therapeutic utility, the phosphotyrosine mimics of this invention may also be used to identify, isolate and purify regulatory proteins or fragments thereof that possess native phosphotyrosine-containing ligands or receptors. For example, it is possible to covalently link the phosphotyrosine mimics according to this invention (or compounds comprising them) to a solid support material, thereby creating an

affinity chromatography matrix. Cell preparations may be passed through such a matrix to identify new tyrosine-kinase dependent regulatory proteins or alternatively, to isolate and purify known proteins. In many cases, phosphotyrosine (or a compound containing a phosphotyrosine residue) cannot be used for this purpose, given
5 phosphotyrosine's strong polarity and tendency to hydrolyze. It will be clear to those of ordinary skill in the art that the generality and multiple utilities of the methods and kits described herein are unique and desirable features of this invention.

The following examples are provided to illustrate the invention described herein.
10 These examples demonstrate various preferred embodiments of this invention and are not to be construed as limiting the scope of the invention in any way.

Examples

15


Example 1

ELISA for the detection of phosphotyrosine mimics

A 96 well microtiter plate (Pierce #15126, Reacti-Bind Streptavidin coated polystyrene) was coated with 100 μ l of biotinylated peptide ECQ(pY)QPQP
20 (MW=1367.5) at a concentration of 15 nM in Tris Buffered Saline (TBS) (10 mM Tris•HCl, pH 7.5, 100 mM NaCl, 0.1% Tween 20) for 45 minutes at room temperature. The microtiter plate was washed 4 times with TBS. MAb PY-20 (Zymed #03-7799) was pre incubated for 45 minutes at a 2X concentration of 0.078 μ g/ml (0.48 nM) with a 2X concentration of non-phosphorylated test compounds in
25 TBS. 100 μ l of test compound/antibody solution was added in duplicate to each of the wells and incubated at room temperature. The microtiter plate was washed 4 times with TBS. 100 μ l of goat anti-mouse IgG (Heavy & Light chains F(ab')₂HRP) was added and diluted to 1:2000 in TBS. After incubating for 45 minutes at room temperature, the well were washed 4 times with TBS. 100 μ l of
30 enzyme substrate (Zymed #00-2011, ABTS Kit) was added as a buffer (9 parts dH₂O to 1 part 10X substrate buffer and 4 drops of 50X ABTS solution added just before use). After 20 minutes, the reaction was stopped by addition of 50 μ l of stop solution. Optical density was measured at 405 nm. Phosphotyrosine mimics were detected by
35 assay. 50% or greater inhibition of anti-phosphotyrosine binding 15 μ g/ml in the above

Example 2Synthesis of N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine
A Compound Containing A Functional Group-Protected Phosphotyrosine Mimic

5



10 A. N-Boc-(para-tributylstannyl)phenylalanine-benzyl ester

A mixture of N-Boc-para-iodophenylalanine-benzyl ester (4.83g), hexabutylditin (6.4 mL) and Pd(Ph₃P)₂Cl₂ (0.101g) in dry toluene (70 mL) was heated at 60 °C under Argon for 55 hours. Aqueous potassium fluoride (20 mL, 1M solution) and tetrabutylammonium fluoride (1M in tetrahydrofuran, 1 mL) were added and the mixture was stirred at room temperature for 1 day. The mixture was diluted with ethyl acetate, washed with water, dried filtered and evaporated. The residue was fractionated over silica gel to give the N-Boc-(para-tributylstannyl)phenylalanine-benzyl ester (3.55 g).

20 B. N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine-benzyl ester.


A mixture of N-Boc-(para-tributylstannyl)phenylalanine-benzyl ester (0.661g), 2-iodobenzoic acid ^tbutyl ester (0.334g) and Pd(Ph₃P)₂Cl₂ (0.035 g) in N-methylpyrrolidone (3 mL) was heated at 110°C for 3 hours under argon. The mixture was diluted with ethyl acetate, washed with water, dried filtered and evaporated. The residue was fractionated over silica gel to give N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine-benzyl ester (0.203 g).

C. N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine.

A mixture of N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine-benzyl ester and 10% Palladium on charcoal (0.030g) in ethanol 20 mL was hydrogenated at 35 psi for 23.5 hours. The reaction mixture was filtered through celite and evaporated to dryness to give the N-Boc-para-(2'-^tbutoxycarbonylphenyl)phenylalanine (0.175 g).

Example 3Representative Synthesis of Phosphotyrosine Mimic-Containing Peptide (BIRE0567)

5



BIRE0567

Peptide synthesis was carried out by solid-phase methodology using Fmoc-chemistry.

10 The side chain protecting group was *tert*-butyl ester for Glu. The solid-phase used was Fmoc-L-Ile-HMPA resin (0.5mmole/g). Coupling was carried out using 3 equivalents of the amino acid using the protocol amino acid: HBTU: HOBT: DIEA-1:1:1:2 molar ratios. Coupling times were typically 45 min. The same protocol was used to couple N-Boc-para-(2'-*t*-butoxycarbonylphenyl)phenylalanine.. Cleavage of the peptide from

15 the resin was achieved by treating the dried resin with trifluoroacetic acid: anisole: ethanedithiol: ethyl methyl sulfide - 95:3:1:1 for 2 hrs at room temperature. The resin was filtered and washed with the cleavage cocktail. After evaporating trifluoroacetic acid under vacuum, the residue was treated with the cold diethyl ether. The precipitated solid was centrifuged off and the solid pellet obtained washed with ether,

20 dissolved in aqueous acetonitrile and lyophilized. N-terminal acetylation was carried out on the crude peptide in water at 0°C for 45 min. using 5 equivalents of acetic anhydride. DIEA was to start the reaction at pH 7. Acetylated peptide was purified by reverse phase hplc on a Vydac C-18 preparative column (300 Å pore size, 10mm particle size; 2.2cm X 25cm) with UV monitoring at 215nm and flow rate of

25 15ml/min. The gradient conditions were 20-100%B over 25min. Mobile phase A was 0.05% TFA/H₂O and B 60% acetonitrile/water containing 0.05% TFA. The major peak was collected and characterized by FABMS to give the desired M+.

While we have herein before presented a number of embodiments of this invention, it is apparent that our basic construction can be altered to provide other embodiments that utilize the processes and compositions of this invention. For example, obvious variations of the synthetic process steps and intermediates described and exemplified
5 herein will be easily recognized by those of ordinary skill in the art. This application expressly envisions and extends to those obvious variations. It should be appreciated that the scope of this invention is defined by the following claims rather than by the specific embodiments that have been presented hereinabove by way of example.

WE CLAIM:

1. A method for identifying phosphotyrosine mimics comprising the steps of:
(a) contacting a non-phosphorylated test compound and a phosphotyrosine containing substance with an anti-phosphotyrosine antibody; and
(b) detecting the ability of the non-phosphorylated test compound to inhibit the binding of the anti-phosphotyrosine antibody to the phosphotyrosine-containing substance.
2. The method according to claim 1, wherein said method comprises the steps of:
(a) competitively binding a non-phosphorylated test compound and a phosphotyrosine-containing substance with an anti-phosphotyrosine antibody; and
(b) detecting the degree to which the non-phosphorylated test compound displaces the phosphotyrosine-containing substance.
3. The method according to claim 2, wherein said method comprises the steps of:
(a) incubating a non-phosphorylated test compound with an anti-phosphotyrosine antibody to form a solution;
(b) contacting the solution of step (a) with a known amount of a phosphotyrosine-containing substance; and
(c) detecting the degree to which the non-phosphorylated test compound displaces the phosphotyrosine-containing substance.
4. The method according to claim 1, wherein said method is an ELISA.
5. The method according to claim 4, wherein said anti-phosphotyrosine antibody is a monoclonal antibody.
6. The method according to claim 5, wherein said monoclonal antibody is PY-20.

7. A phosphotyrosine mimic identified by the method according to claim 1.
8. A phosphotyrosine mimic according to claim 7 wherein said mimic is phenyl carboxylic acid.
9. A method for inhibiting the activation of a phosphotyrosine dependent regulatory protein comprising the steps of:
 - (a) incorporating a phosphotyrosine mimic according to claim 7 or 8 into an appropriate molecular structure to produce a phosphotyrosine mimic-containing compound and
 - (b) contacting the compound of step (a) with a tyrosine kinase-dependent regulatory protein.
10. The method according to claim 9, wherein the compound of step (a) is covalently attached to a solid support prior to the binding step (b).
11. A kit for identifying phosphotyrosine mimics comprising:
 - (a) an immobilized phosphotyrosine-containing substance and
 - (b) an anti-phosphotyrosine antibody, wherein said antibody is conjugated to a detectable label.
12. A kit for identifying phosphotyrosine mimics comprising:
 - (a) an immobilized anti-phosphotyrosine antibody and
 - (b) a phosphotyrosine-containing substance, wherein said substance is conjugated to a detectable label.
13. A phosphotyrosine mimic which is a phenyl carboxylic acid or a functional group protected derivative thereof.

14. A method for treating a neoplastic or chronic inflammatory disease comprising the step of administering to a patient a therapeutically effective amount of a phosphotyrosine mimic-containing peptidomimetic of a phosphotyrosine-containing ligand or receptor of a tyrosine-kinase dependent regulatory protein.
15. The method according to claim 14, wherein the phosphotyrosine mimic is a phenyl carboxylic acid or a functional group protected derivative thereof.

INTERNATIONAL SEARCH REPORT

Intern. Application No. **PCT/US 97/07127**

A. CLASSIFICATION OF SUBJECT MATTER

G 01 N 33/53, G 01 N 33/577, G 01 N 33/543, A 61 K 35/00,
A 61 K 31/19, A 61 K 31/195, A 61 K 31/66

According to International Patent Classification (IPC) or to both national classification and IPC⁶

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G 01 N, A 61 K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, A, 93/03 377 (ZILTENER) 18 February 1993 (18.02.93), abstract; claims. --	1-6
A	EP, A, 0 449 269 (BOEHRINGER MANNHEIM GMBH) 02 October 1991 (02.10.91), abstract; claims. --	1-6
A	DATABASE WPIL, AN 93-201 126, DERWENT PUBLICATIONS LTD., London; & JP, A, 05 126 833 (HIRANO, TOSOH COR), abstract. --	1-6
A, P	WO, A, 97/07 131 --	7, 8,

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search
14 August 1997

Date of mailing of the international search report

11. 09. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HX Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

SCHNASS e.h.

INTERNATIONAL SEARCH REPORT

-2-

Internat'l Application No

PCT/US 97/07127

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	(CIBA-GEIGY AG) 27 February 1997 (27.02.97), claims.	13-15
A	-- CHEMICAL ABSTRACTS, vol. 123, no. 17, issued 1995, October 23, (Columbus, Ohio, USA), YE BIN et al. "L-O-(2-Malo- nyl)tyrosine: A New Phospho- tyrosyl Mimetic for the Pre- paration of Src Homology 2 Domain Inhibitory Peptides", page 1265, column 1, no. 228 860z; & J. Med. Chem. 1995, 38(21), 4270-5 (Eng).	7,8,13
A	-- CHEMICAL ABSTRACTS, vol. 123, no. 1, issued 1995, July 03, (Columbus, Ohio, USA), H.K. KOLE et al. "Protein- -tyrosine phosphatase inhibi- tion by a peptide containing the phosphotyrosyl mimetic, L-O-malonyltyrosine", pages 449-450, no. 4 299k; & Biochem. Biophys. Res. Commun. 1995, 209(3), 817-22 (Eng).	7,8,13
A,P	-- CHEMICAL ABSTRACTS, vol. 125, no. 13, issued 1996, September 23, (Columbus, Ohio, USA), A. SHAHRIPOUR et al. "Novel phosphotyrosine mimetics in the design of peptide ligands for pp60srcSH2 domain", pages 1230-1231, no. 168 619g; & Bioorg. Med. Chem. Lett. 1996, 6(11), 1209-1214 (Eng).	7,8,13
A,P	-- CHEMICAL ABSTRACTS, vol. 125, no. 7, issued 1996, August 12, (Columbus, Ohio, USA), A.K. KZARDENINGS et al. "A general and convenient syn- thesis of novel phospho- tyrosine mimetics", page 1236, column 2,	7,8, 13

INTERNATIONAL SEARCH REPORT

Internat Application No

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT			PCT/US 97/07127
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
A	<p>no. 87 136v; & Tetrahedron Lett. 1996, 37(21), 3635-3638 (Eng). --</p> <p>WO, A, 95/25 118 (TRUSTEES OF TUFTS UNIVER- SITY) 21 September 1995 (21.09.95), claims 1-30, 44-46. ----</p>	7, 8, 13	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/07127

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Though the subject-matter of claims 14 and 15 is excluded from patentability by rule 39.1(iv)PCT it was searched, too.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/US 97/07127 SAE 160394

In diesem Anhang sind die Mitglieder
der Patentfamilien der im obenge-
nannten internationalen Recherchenbericht
angeführten Patentdokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visée ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
WO A1 9303377	18-02-93	keine - none - rien	
EP A1 449269	02-10-91	AT E 138205 DE A1 4009848 DE C0 59107797 EP B1 449269 ES T3 2089048 JP A2 4223270 JP B4 7104347 US A 5580742	15-06-96 02-10-91 20-06-96 15-05-96 01-10-96 13-08-92 13-11-95 03-12-96
JP A2 5126830	21-05-93	keine - none - rien	
WO A1 9707131	27-02-97	AU A1 68709/96 GB A0 9516842	12-03-97 18-10-95
WO A2 9525118	21-09-95	US A 5580979 WO A3 9525118	03-12-96 14-11-95